# **Purina® Honor® Show Fitter's Edge 13 DX**



Specie: Show Cattle Type of Feed: Complete Form of Feed: Textured

#### **General Description:**

The versatility of Honor® Show Fitter's Edge 13 DX makes this the "tried-and-true" staple of the Honor® Show lineup. Providing flexibility from start to finish, Fitter's Edge® can be fed as a complete ration to grow and develop show cattle.

# Purina® Honor® Show Fitter's Edge® 13 DX MEDICATED

Type C Medicated Feed FOR CATTLE 400 LBS TO SHOW

For the prevention of coccidiosis in cattle (ruminating and non-ruminating calves and veal calves) by Eimeria bovis and E. zuernii.

#### **ACTIVE DRUG INGREDIENTS**

Decoquinate	0.00250%
	(11.35 mg/lb)
GUARANTEED ANALYS	IS
Crude Protein, (Min)	13.00%
Crude Fat, (Min)	2.50%
Crude Fiber, (Max)	
Calcium (Ca), (Min)	
Calcium (Ca), (Max)	1.08%
Phosphorus (P), (Min)	
Salt (NaCl), (Min)	
Salt (NaCl), (Max)	0.65%
Sodium (Na), (Max)	
Potassium (K), (Min)	0.65%
Selenium (Se), (Min)	
Selenium (Se), (Max)	
Vitamin A, (Min)	

#### \*INGREDIENTS

Flaked Corn, Cottonseed Hulls, Whole Oats, Dehulled Soybean Meal, Wheat Middlings, Cane Molasses, Cracked Corn, Soybean Oil, Calcium Carbonate, Salt, Monocalcium Phosphate, Dicalcium Phosphate, Propionic Acid (a Preservative), Active Dry Yeast, Molasses Hydrolyzed Yeast, Vitamin A Supplement, Acetic Acid, Ammonium Hydroxide, Benzoic Acid (a Preservative), Natural Flavor, Artificial Flavor, Sorbic Acid (a Preservative), Vitamin D3 Supplement, Methylparaben (a Preservative), Organic Soybean Oil, Silicon Dioxide, Zinc Amino Acid Complex, Manganese Amino Acid Complex, Copper Amino Acid Complex, Cobalt Glucoheptonate, Propylparaben (a Preservative), Rice Hulls, Saccharin Sodium, Calcium Chloride, Vitamin E Supplement, Cobalt Carbonate, Manganese Sulfate, Ethylenediamine Dihydroiodide, Zinc Sulfate, Basic Copper Chloride, Sodium Selenite

\*Note: ingredients differ by manufacturing plant. Please refer to the tag for a current list of ingredients.

#### FEEDING DIRECTIONS:

Feed at the rate of 2 lbs per 100 lbs of bodyweight to provide 22.7 mg Decoquinate per 100 pounds of body weight (0.5 mg/kg) per day. Feed at least 28 days during period of exposure to coccidiosis or when it is likely to be a hazard.

Follow these management practices:

- Monitor body condition of cattle on a regular basis.
- Recommended feeding rates should be adjusted depending on cattle body condition.
- When making a ration change, allow 7 to 10 days for animals to adjust to the new ration.
- Provide adequate feeder space for each animal.
- Provide mineral and salt free choice.
- Provide fresh, clean water near the feeding area. In groups of cattle, there are certain animals that experience chronic bloat or other digestive disturbances and consequently, are poor performers. In addition, excess feed consumption, severe weather changes resulting in erratic feed consumption and poorly managed feeding practices can increase the incidence of bloat in all cattle. If bloat occurs, please review feed management practices.

## Available Options\*:



Product No. / Formula Code	Package	Form/Size Diameter
3011976-506 / 39ZS	Bag 50#	Textured – Flaked Corn
3011975-506 / 39ZY	Bag 50#	Textured - Cracked Corn

<sup>\*</sup>Product version depends on regional availability

Product Features:	Product Benefits:
A uniquely-formulated versatile ration	<ul> <li>Offers flexibility from start to finish in your feeding program</li> <li>Can be fed as a complete ration to grow and develop show cattle</li> <li>Blend with High Octane® Power Fuel® to create an outstanding finishing ration</li> </ul>
Highly palatable	Hand-picked ingredients proven to promote intake and keep cattle on feed during the stresses of show season
Peloton® Yeast	<ul> <li>Thermal stable yeast encapsulation maintains effectiveness of yeapelleting.</li> <li>Optimized feed digestibility, supporting feed intake, and animal performance from Saccharomyces cerevisiae yeast.</li> <li>PMI Fulfill® Feed Additive Mannans (MOS) and beta-glucans sugut integrity and immune-competence.</li> </ul>
Zinpro Availa® 4 Minerals	<ul> <li>Balanced combination of zinc, manganese, copper, and cobalt</li> <li>Provides critical, highly bioavailable minerals that help support is function and hoof integrity.</li> </ul>
Medicated with Deccox® (decoquinate)	For the prevention of coccidiosis

#### WARNING:

Do not feed to cows producing milk for human consumption.

### CAUTION:

Bentonite should not be used in decoquinate feeds. DO NOT FEED TO SHEEP DUE TO HIGH LEVELS OF SUPPLEMENTAL COPPER. Follow label directions: Feeding added selenium at levels in excess of 0.3 ppm in total diet is prohibited. Store in a dry, well-ventilated area protected from rodents and insects. Do not feed moldy or insect-infested feed to animals as it may cause illness, performance loss or death. USE ONLY AS DIRECTED