

## 567/1098 BUTLER BEEF MIX

# MEDICATED FOR CATTLE ON PASTURE

Pasture cattle (slaughter, stocker, feeder cattle, and dairy and beef replacement heifers): For increased rate of weight gain. Intakes of lasalocid in excess of 200 mg/head/day have not been shown to be more effective than 200 mg/head/day.

### **ACTIVE DRUG INGREDIENT**

Lasalocid	20 g/ton
-----------	----------

GUARANTEED ANALYSIS						
Crude Protein	(Min)	10.00 %	<b>Phosphorus</b>	(Min)	0.30 %	
Crude Fat	(Min)	3.70 %	Salt	(Min)	0.50 %	
Crude Fiber	(Max)	14.00 %	Salt	(Max)	0.90 %	
Calcium	(Min)	0.60 %	Potassium	(Min)	0.90 %	
Calcium	(Max)	1.00 %	Vitamin A	(Min)	8,000 IU/LB	

#### **INGREDIENTS**

Grain Products, Roughage Products (20%), Forage Products, Molasses Products, Plant Protein Products, Processed Grain Byproducts, Ground Limestone, Soybean Oil, Salt, Selenium Yeast, Cobalt Carbonate, Copper Chloride, Copper Sulfate, Ethylenediamine Dihydriodide, Manganese Sulfate, Manganous Oxide, Zinc Sulfate, Zinc Oxide, Mineral Oil, Calcium Carbonate, Vitamin E Supplement, Vitamin A Supplement and Vitamin D3 Supplement.

#### FEEDING DIRECTIONS

Feed this product at a rate of 6 to 30 pounds per head per day to provide no less than 60 milligrams nor more than 300 milligrams of lasalocid per head per day. Intake of lasalocid in excess of 200 mg/head/day has not been shown to be more effective than 200 mg/head/day. Always provide adequate amounts of good forage (hay or pasture) and plenty of fresh, clean water.

**WARNING**: A withdrawal period has not been established for this product in pre-ruminating calves. Do not use in calves to be processed for veal.

**CAUTION:** The safety of Lasalocid for use in unapproved species has not been established. Do not allow horses or other equines access to feeds containing lasalocid as ingestion may be fatal.

Manufactured by PRODUCERS COOPERATIVE ASSOCIATION Bryan, Texas 77806

Net Weight 50 lbs. (22.68 kg) or Bulk