

**LINDNER**  
**632 Starter / Grower BMD-FBZ**  
**MEDICATED**

A complete feed for Starter/Grower pigs

Bacitracin Methylene Disalicylate is indicated for control of swine dysentery associated with *Treponema hyodysenteriae* on the premises with a history of swine dysentery but where signs of the disease have not yet occurred, or following an approved treatment of the disease condition.

Fenbendazole is indicated for the removal of: Adult stage Lungworms: *Metastrongylus apri*, *Metastrongylus pudendotectus*. Gastrointestinal worms: Adult and larvae (L3, L4 stages, liver, lung, intestinal forms) large roundworm (*Ascaris suum*); adult stage nodular worms (*Oesophagostomum dentatum*, *Ornithostrongylus quadrispinulatum*); adult stage small stomach worm (*Hyostromylus rubidus*); Adult and larvae (L2, L3, L4 stages- intestinal mucosal forms) whipworm (*Trichuris suis*); Kidneyworm: Adult and larvae *Stephanurus dentatus*.

**ACTIVE DRUG INGREDIENTS**

Bacitracin Methylene Disalicylate ..... 250 g/ton  
 Fenbendazole ..... 29.2 g/ton(14.6 mg/lb)

**GUARANTEED ANALYSIS**

Crude Protein (Min) ..... 20.0%  
 Lysine (Min) ..... 1.4%  
 Crude Fat (Min) ..... 5.0%  
 Crude Fiber (Max) ..... 3.0%  
 Calcium (Min) ..... 0.7%  
 Calcium (Max) ..... 1.2%  
 Phosphorous (Min) ..... 0.7%  
 Salt (Min) ..... 0.6%  
 Salt (Max) ..... 1.1%  
 Selenium (Min) ..... 0.3 ppm  
 Zinc (Min) ..... 120 ppm

**INGREDIENTS**

Grain Products, Plant Protein Products, Animal Protein Products, Animal Fat, Monocalcium Phosphate, Calcium Carbonate, Sodium Bentonite, Salt, Roughage Products, Ferrous Sulfate, Zinc Sulfate, Choline Chloride, Vitamin E Supplement, Copper Sulfate, Dried *Bacillus Licheniformis* Fermentation Product, Dried *Bacillus Subtilis* Fermentation Product, Sodium Aluminosilicate, D-Calcium Pantothenate, Manganese Sulfate, Niacin Supplement, Mineral Oil, Vitamin B12 Supplement, Dried *Trichoderma Reesei* Fermentation Products, Sodium Sulfate, Hydrated Magnesium Silicate, Poly-Vinyl Alcohol, Riboflavin Supplement, Sodium Selenite, Vitamin A Supplement, Menadione Sodium Bisulfite Complex, Vitamin D3 Supplement, Ethylenediamine Dihydriodide, DL-Methionine, L-Lysine, L-Threonine, Artificial Flavor

**FEEDING DIRECTIONS**

Feed as the sole ration to provide a total intake of 9mg Fenbendazole/kg of body weight (4.08mg Fenbendazole/lb. of body weight) divided over 3-12 days. Multiply 0.28 times pound of body weight and feed this amount of product over 3-12 days.

Example:

<u>Pig to be Dewormed</u>	<u>Starter/Grower Feed Required per pig</u>	<u>Days Required to Consume Dose</u>
50 lb.	14.0 lb.	7
75 lb.	21.0 lb.	7
100 lb.	28.0 lb.	7

**CAUTION:** Not for use in swine that weigh more than 250 lb. Under conditions of continued exposure to parasites, retreatment may be needed after 4-6 weeks. Diagnosis of swine dysentery should be confirmed by a veterinarian when results are not satisfactory.

**CONSULT YOUR VETERINARIAN FOR ASSISTANCE IN THE DIAGNOSIS, TREATMENT, AND CONTROL OF PARASITISM.**

**WARNING:** Parasite resistance may develop to any dewormer. All dewormers require accurate dosing for best results. Following the use of any dewormer, effectiveness of treatment should be monitored. A decrease of effectiveness over time may indicate the development of resistance to the dewormer administered. The parasite management plan should be adjusted accordingly based on regular monitoring.

**WITHDRAWAL PERIODS**

**Swine must not be slaughtered for human consumption within 4 days following last treatment with this drug product.**

**MANUFACTURED BY**

LINDNER FEED AND MILLING CO., INC  
 Comfort, Texas 78013  
 Net Weight 50 lb (22.68 kg)

**LINDNER**  
**632 Starter / Grower BMD-FBZ**  
**MEDICATED**

A complete feed for Starter/Grower pigs

Bacitracin Methylene Disalicylate is indicated for control of swine dysentery associated with *Treponema hyodysenteriae* on the premises with a history of swine dysentery but where signs of the disease have not yet occurred, or following an approved treatment of the disease condition.

Fenbendazole is indicated for the removal of: Adult stage Lungworms: *Metastrongylus apri*, *Metastrongylus pudendotectus*. Gastrointestinal worms: Adult and larvae (L3, L4 stages, liver, lung, intestinal forms) large roundworm (*Ascaris suum*); adult stage nodular worms (*Oesophagostomum dentatum*, *Ornithostrongylus quadrispinulatum*); adult stage small stomach worm (*Hyostromylus rubidus*); Adult and larvae (L2, L3, L4 stages- intestinal mucosal forms) whipworm (*Trichuris suis*); Kidneyworm: Adult and larvae *Stephanurus dentatus*.

**ACTIVE DRUG INGREDIENTS**

Bacitracin Methylene Disalicylate ..... 250 g/ton  
 Fenbendazole ..... 29.2 g/ton(14.6 mg/lb)

**GUARANTEED ANALYSIS**

Crude Protein (Min) ..... 20.0%  
 Lysine (Min) ..... 1.4%  
 Crude Fat (Min) ..... 5.0%  
 Crude Fiber (Max) ..... 3.0%  
 Calcium (Min) ..... 0.7%  
 Calcium (Max) ..... 1.2%  
 Phosphorous (Min) ..... 0.7%  
 Salt (Min) ..... 0.6%  
 Salt (Max) ..... 1.1%  
 Selenium (Min) ..... 0.3 ppm  
 Zinc (Min) ..... 120 ppm

**INGREDIENTS**

Grain Products, Plant Protein Products, Animal Protein Products, Animal Fat, Monocalcium Phosphate, Calcium Carbonate, Sodium Bentonite, Salt, Roughage Products, Ferrous Sulfate, Zinc Sulfate, Choline Chloride, Vitamin E Supplement, Copper Sulfate, Dried *Bacillus Licheniformis* Fermentation Product, Dried *Bacillus Subtilis* Fermentation Product, Sodium Aluminosilicate, D-Calcium Pantothenate, Manganese Sulfate, Niacin Supplement, Mineral Oil, Vitamin B12 Supplement, Dried *Trichoderma Reesei* Fermentation Products, Sodium Sulfate, Hydrated Magnesium Silicate, Poly-Vinyl Alcohol, Riboflavin Supplement, Sodium Selenite, Vitamin A Supplement, Menadione Sodium Bisulfite Complex, Vitamin D3 Supplement, Ethylenediamine Dihydriodide, DL-Methionine, L-Lysine, L-Threonine, Artificial Flavor

**FEEDING DIRECTIONS**

Feed as the sole ration to provide a total intake of 9mg Fenbendazole/kg of body weight (4.08mg Fenbendazole/lb. of body weight) divided over 3-12 days. Multiply 0.28 times pound of body weight and feed this amount of product over 3-12 days.

Example:

<u>Pig to be Dewormed</u>	<u>Starter/Grower Feed Required per pig</u>	<u>Days Required to Consume Dose</u>
50 lb.	14.0 lb.	7
75 lb.	21.0 lb.	7
100 lb.	28.0 lb.	7

**CAUTION:** Not for use in swine that weigh more than 250 lb. Under conditions of continued exposure to parasites, retreatment may be needed after 4-6 weeks. Diagnosis of swine dysentery should be confirmed by a veterinarian when results are not satisfactory.

**CONSULT YOUR VETERINARIAN FOR ASSISTANCE IN THE DIAGNOSIS, TREATMENT, AND CONTROL OF PARASITISM.**

**WARNING:** Parasite resistance may develop to any dewormer. All dewormers require accurate dosing for best results. Following the use of any dewormer, effectiveness of treatment should be monitored. A decrease of effectiveness over time may indicate the development of resistance to the dewormer administered. The parasite management plan should be adjusted accordingly based on regular monitoring.

**WITHDRAWAL PERIODS**

**Swine must not be slaughtered for human consumption within 4 days following last treatment with this drug product.**

**MANUFACTURED BY**

LINDNER FEED AND MILLING CO., INC  
 Comfort, Texas 78013  
 Net Weight 50 lb (22.68 kg)